


How to use

Prenoxad[®] Naloxone Injection

as part of an opioid overdose response



Step 1

Safety


Check the person and area for anything unsafe.



Step 2

Opioid overdose? Look for the signs

- Not responding
- Pinned pupils
- Not breathing normally - shallow slow breathing or snoring
- Pale or cold skin, blue or grey lips and fingertips



Step 3


**Call Triple Zero (000)
and put on speaker phone while you
administer Prenoxad[®]**



Step 4


Lie person on side

Put person in recovery position.



Step 5

Screw needle onto Prenoxad® prefilled syringe




Step 6

Position the needle

Insert needle into outer thigh or upper arm.


Do not inject into a vein.
(You do not need to remove clothing first).



Step 7

Give one dose of naloxone

Inject 0.4mL, up to 1st black line on syringe (Prenoxad® syringe contains 5 doses).



Step 8

Are they breathing?

If not, use rescue breathing and/or chest compressions.


(2 breaths / 30 compressions)



Step 9

Recovery

Put person in recovery position. Stay with them until ambulance arrives.



Step 10

Repeat dose if needed

If no response inject another 0.4mL dose (to next black line) every 2 minutes until person recovers or ambulance arrives.



Prenoxad® takes 2-5 minutes to start working and lasts 30-90 minutes.

For information and support call The National Alcohol and Drug Hotline, 24 hours a day, 7 days a week on **1800 050 015** for free from anywhere in Australia.